

# CONTENTS OF No. 532, April 2017

Page

LEONE MELLUSO, VINCENZA GUARINO, MICHELE LUSTRINO, VINCENZO MORRA and ROBERTO DE' GENNARO: The <i>REE</i> - and HFSE-bearing phases in the Itatiaia alkaline complex (Brazil) and geochemical evolution of feldspar-rich felsic melts	<b>217</b>
M. LACALAMITA, E. MESTO, E. KANEVA, F. SCORDARI, G. PEDRAZZI, N. VLADYKIN and E. SCHINGARO: Structure refinement and crystal chemistry of tokkoite and tinaksite from the Murun massif (Russia)	<b>251</b>
ANTHONY R. KAMPE, JAKUB PLÁŠIL, ANATOLY V. KASATKIN, JOE MARTY, JIŘÍ ČEJKA and LADISLAV LAPČÁK: Shumwayite, [(UO <sub>2</sub> )(SO <sub>4</sub> )(H <sub>2</sub> O) <sub>2</sub> ] <sub>2</sub> ·H <sub>2</sub> O, a new uranyl sulfate mineral from Red Canyon, San Juan County, Utah, USA	<b>273</b>
I. E. GREY, E. KECK, A. R. KAMPE, C. M. MACRAE, A. M. GLENN and J. R. PRICE: Wilhelmgümbelite, [ZnFe <sup>2+</sup> Fe <sup>3+</sup> (PO <sub>4</sub> ) <sub>3</sub> (OH) <sub>4</sub> (H <sub>2</sub> O) <sub>5</sub> ] <sub>2</sub> ·2H <sub>2</sub> O, a new schoonerite-related mineral from the Hagendorf Süd pegmatite, Bavaria	<b>287</b>
MARK D. WELCH and ANTHONY R. KAMPE: Stoichiometric partially-protonated states in hydroxide perovskites: the jeanbandyite enigma revisited	<b>297</b>
F. CÁMARA, E. BITTARELLO, M. E. CIRIOTTI, F. NESTOLA, F. RADICA, F. MASSIMI, C. BALESTRA and R. BRACCO: As-bearing new mineral species from Valletta mine, Maira Valley, Piedmont, Italy: III. Canosioite, Ba <sub>2</sub> Fe <sup>3+</sup> (AsO <sub>4</sub> ) <sub>2</sub> (OH), description and crystal structure	<b>305</b>
ANTHONY R. KAMPE, BARBARA P. NASH, JOE MARTY and JOHN M. HUGHES: Mesaite, CaMn <sup>2+</sup> (V <sub>2</sub> O <sub>7</sub> ) <sub>3</sub> ·12H <sub>2</sub> O, a new vanadate mineral from the Packrat mine, near Gateway, Mesa County, Colorado, USA	<b>319</b>
I. E. GREY, E. KECK, A. R. KAMPE, W. G. MUMME, C. M. MACRAE, R. W. GABLE, A. M. GLENN and C. J. DAVIDSON: Steinmetzite, Zn <sub>2</sub> Fe <sup>3+</sup> (PO <sub>4</sub> ) <sub>2</sub> (OH)·3H <sub>2</sub> O, a new mineral formed from alteration of phosphophyllite at the Hagendorf Süd pegmatite, Bavaria	<b>329</b>
E. SCHINGARO, E. MESTO, M. LACALAMITA, F. SCORDARI, E. KANEVA and F. N. VLADYKIN: Single-crystal X-ray diffraction, EMPA, FTIR and X-ray photoelectron spectroscopy study of narsarsukite from Murun Massif, Russia	<b>339</b>
DAN TOPA and EMIL MAKOVICKY: The crystal structure of veenite	<b>355</b>
F. CÁMARA, E. SOKOLOVA, Y. A. ABDU, F. C. HAWTHORNE, T. CHARRIER, V. DORCET and J. -F. CARPENTIER: Fogoite-(Y), Na <sub>3</sub> Ca <sub>2</sub> Y <sub>2</sub> Ti(Si <sub>2</sub> O <sub>7</sub> ) <sub>2</sub> OF <sub>3</sub> , a Group I TS-block mineral from the Lagoa do Fogo, the Fogo volcano, São Miguel Island, the Azores: Description and crystal structure	<b>369</b>
A. N. SAPOZHNIKOV, E. V. KANEVA, L. F. SUVOROVA, V. I. LEVITSKY and L. A. IVANOVA: Sulfhydrylbystrite, Na <sub>5</sub> K <sub>2</sub> Ca(Al <sub>6</sub> Si <sub>6</sub> O <sub>24</sub> )(S <sub>5</sub> )(SH), a new mineral with the LOS framework, and re-interpretation of bystrite: cancrinite-group minerals with novel extra-framework anions	<b>383</b>
<i>CNMNC Newsletter 36</i>	
U. HÄLENIUS, F. HATERT, M. PASERO and S.J. MILLS: New minerals and nomenclature modifications approved in 2017	<b>403</b>